


Zimbabwe

Epidemiological Fact Sheets

on HIV/AIDS
and Sexually
Transmitted
Infections



2002 Update



Estimated number of people living with HIV/AIDS

In 2001 and during the first quarter of 2002, UNAIDS and WHO worked closely with national governments and research institutions to recalculate current estimates on people living with HIV/AIDS. These calculations are based on the previously published estimates for 1997 and 1999 and recent trends in HIV/AIDS surveillance in various populations. A methodology developed in collaboration with an international group of experts was used to calculate the new estimates on prevalence and incidence of HIV and AIDS deaths, as well as the number of children infected through mother-to-child transmission of HIV. Different approaches were used to estimate HIV prevalence in countries with low-level, concentrated or generalized epidemics. The current estimates do not claim to be an exact count of infections. Rather, they use a methodology that has thus far proved accurate in producing estimates which give a good indication of the magnitude of the epidemic in individual countries. However, these estimates are constantly being revised as countries improve their surveillance systems and collect more information.

Adults in this report are defined as women and men aged 15 to 49. This age range covers people in their most sexually active years. While the risk of HIV infection obviously continues beyond the age of 50, the vast majority of those who engage in substantial risk behaviours are likely to be infected by this age. The 15 to 49 range was used as the denominator in calculating adult HIV prevalence.

■ Estimated number of adults and children living with HIV/AIDS, end of 2001

These estimates include all people with HIV infection, whether or not they have developed symptoms of AIDS, alive at the end of 2001:

Adults and children	2,300,000		
Adults (15-49)	2,000,000	Adult rate (%)	33.7
Women (15-49)	1,200,000		
Children (0-15)	240,000		

■ Estimated number of deaths due to AIDS

Estimated number of adults and children who died of AIDS during 2001:

Deaths in 2001	200,000
-----------------------	----------------

■ Estimated number of orphans

Estimated number of children who have lost their mother or father or both parents to AIDS and who were alive and under age 15 at the end of 2001:

Current living orphans	780,000
-------------------------------	----------------

UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance

Global Surveillance of HIV/AIDS and sexually transmitted infections (STIs) is a joint effort of WHO and UNAIDS. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, initiated in November 1996, guides respective activities. The primary objective of the Working Group is to strengthen national, regional and global structures and networks for improved monitoring and surveillance of HIV/AIDS and STIs. For this purpose, the Working Group collaborates closely with national AIDS programmes and a number of national and international experts and institutions. The goal of this collaboration is to compile the best information available and to improve the quality of data needed for informed decision-making and planning at national, regional, and global levels. The Epidemiological Fact Sheets are one of the products of this close and fruitful collaboration across the globe.

Within this framework, the Fact Sheets collate the most recent country-specific data on HIV/AIDS prevalence and incidence, together with information on behaviours (e.g. casual sex and condom use) which can spur or stem the transmission of HIV.

Not unexpectedly, information on all of the agreed-upon indicators was not available for many countries in 2001. However, these updated Fact Sheets do contain a wealth of information which allows identification of strengths in currently existing programmes and comparisons between countries and regions. The Fact Sheets may also be instrumental in identifying potential partners when planning and implementing improved surveillance systems.

The fact sheets can be only as good as information made available to the UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance. Therefore, the Working Group would like to encourage all programme managers as well as national and international experts to communicate additional information to them whenever such information becomes available. The Working Group also welcomes any suggestions for additional indicators or information proven to be useful in national or international decision-making and planning.

Assessment of the epidemiological situation (2002)


HIV prevalence among ANC attendees tested during the 2000 HIV sentinel surveillance survey at 19 sites distributed across the country was 35%; sixteen sites had rates exceeding 25%. HIV infection rates ranged from 25.7% to 70.7%. Comparing the HIV prevalence of 1997 and 2000 at the 10 sites used in both surveys, median HIV prevalence increased from 27.4% to 36.1%. In 2000, ANC attendees aged 15-19 years had an HIV prevalence of 27.8%, the 20-24 year-olds had a rate of 35.1%, and the 30-34 year olds had the highest rate of 43.5%. HIV prevalence was higher among women living in commercial farming areas (43.7%) and in growth points (38.3%) within rural areas. In the major urban areas, HIV prevalence among antenatal clinic attendees tested increased from 10% in 1989 to 36% in 1994. In 1997, 30% of antenatal clinic attendees tested HIV positive. Age detail is available from Harare in 1995 only: 26% of antenatal clinic attendees under 20 years of age tested positive for HIV, including 28% of women 15-17 years of age. Outside of Harare, sentinel surveillance information among antenatal clinic attendees is available since 1990. HIV prevalence among antenatal clinic attendees tested has increased from 12% in 1990 to 37% in 1995; in 1997, a median of 30% of antenatal clinic women tested in 31 sites were HIV positive. In Masvingo in 1995, where 42% of antenatal clinic attendees tested were HIV positive, 34% of women less than 20 years of age were HIV positive. Peak infection in the Masvingo site occurred among women 20 to 24 years of age with 49% testing positive for HIV. ANC surveillance survey was conducted in 2001 and the results will be available around June 2002.

There is only one study available with information on HIV prevalence among sex workers in Zimbabwe. In 1994-95, 86% of sex workers tested in Harare were HIV positive. In Harare, HIV prevalence among STI clinic patients tested increased from 52% in 1990 to 71% in 1995. Outside of Harare, HIV prevalence among STI clinic patients increased from 6% in 1987 to 72% in 1996.

Antenatal clinic women at the 19 sentinel sites were screened for syphilis with Trepanoma Pallidum Heamoagglutination test (TPHA) during the 2000 HIV sentinel surveillance survey. 5% of the women screened had a positive syphilis serology. Women in the age group 35-39 years had the highest rate of 12.8%.

Country Information

Population pyramid, 2001



Indicators	Year	Estimate	Source
Total Population (thousands)	2001	12,852	UNPOP
Population Aged 15-49 (thousands)	2001	5,972	UNPOP
Annual Population Growth	1995-2000	1.9	UNPOP
% of Urban Population	2000	35	UNPOP
Average Annual Growth Rate of Urban Population	1995-2000	3.5	UNPOP
GNI Per Capita (US\$)	1999	530	World Bank
GNI Per Capita Average Annual Growth Rate	1999	-2.6	World Bank
Per Capita Expenditure of Health	1997	49	World Bank
% of Government Budget Spent on Health Care	1998	17.0	WHO
Total Adult Literacy Rate	1997	86	UNESCO
Adult Male Literacy Rate	1997	91	UNESCO
Adult Female Literacy Rate	1997	82	UNESCO
Male Primary School Enrolment Ratio	1996	114.8	UNESCO
Female Primary School Enrolment Ratio	1996	111.3	UNESCO
Male Secondary School Enrolment Ratio	1996	52.4	UNESCO
Female Secondary School Enrolment Ratio	1996	44.6	UNESCO
Crude Birth Rate (births per 1,000 pop.)	1995-2000	37	UNPOP
Crude Death Rate (deaths per 1,000 pop.)	1995-2000	18	UNPOP
Maternal Mortality Rate (per 100,000 live births)	1995	610	WHO
Life Expectancy at Birth	1995-2000	43	UNPOP
Total Fertility Rate	1995-2000	5.0	UNPOP
Infant Mortality Rate (per 1,000 live births)	1995-2000	65	UNPOP
Under 5 Mortality Rate	1995-2000	108	UNPOP

For consistency reasons the data used in the above table are taken from official UN publications.

Contact address:

UNAIDS/WHO Working Group on Global HIV/AIDS
and STI Surveillance
20, Avenue Appia
CH - 1211 Geneva 27
Switzerland
Fax: +41-22-791-4834

email: HIV-AIDS@who.int

<http://www.who.int>
<http://www.unaids.org>

4 - Zimbabwe

HIV prevalence in different populations

This section contains information about HIV prevalence in different populations. The data reported in the tables below are mainly based on the HIV database maintained by the United States Bureau of the Census where data from different sources, including national reports, scientific publications and international conferences are compiled. To provide a simple overview of the current situation and trends over time, summary data are given by population group, geographical area (Major Urban Areas versus Outside Major Urban Areas), and year of survey. Studies conducted in the same year are aggregated and the median prevalence rates (in percentages) are given for each of the categories. The maximum and minimum prevalence rates observed, as well as the total number of surveys/sentinel sites, are provided with the median, to give an overview of the diversity of HIV-prevalence results in a given population within the country. Data by sentinel site or specific study from which the medians were calculated are printed at the end of this fact sheet.

The differentiation between the two geographical areas Major Urban Areas and Outside Major Urban Areas is not based on strict criteria, such as the number of inhabitants. For most countries, Major Urban Areas were considered to be the capital city and - where applicable - other metropolitan areas with similar socio-economic patterns. The term Outside Major Urban Areas considers that most sentinel sites are not located in strictly rural areas, even if they are located in somewhat rural districts.

■ HIV sentinel surveillance

Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
Pregnant women	Major Urban Areas	N-sites			1	4	1		1	2	2	1	6				3
		Minimum			10	16	17.1		25.8	30.4	30	30.4	22.7				30
		Median			10	18.65	17.1		25.8	36.05	31	30.4	30				31.1
	Maximum			10	23.8	17.1		25.8	41.7	32	30.4	34				33.5	
	Outside Major Urban Areas	N-sites			4	14	16		12	19	11	4	30	3	2		18
		Minimum			7.6	7.7	0	13.7	14	19	34	7	29	20.8	13		
Median				16	18.5	0	21.25	24.3	39.5	41.6	28.65	31	22.3	33.15			
Maximum			31.6	33.8	6.6	27	40	70.2	59	53.3	37	23.8	70.7				

Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
Sex workers	Major Urban Areas	N-sites										1					
		Minimum										86					
		Median										86					
		Maximum										86					

Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
Injecting drug users																	

Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
STI patients, Males/both & females	Major Urban Areas	N-sites				1	2	2	1		3	1					
		Minimum				52	39	56	60.3		50.7	53					
		Median				52	45	63	60.3		71	53					
	Maximum				52	51	70	60.3		71.2	53						
	Outside Major Urban Areas	N-sites	2		4	15	15	9	8	7	1						
		Minimum	5.2		24.3	24.49	39	25.6	48	43	71.8						
Median		5.9		30.65	45.6	44.6	52	54	65	71.8							
Maximum	6.6		59.7	65.3	59.6	71.8	75.4	87.7	71.8								

Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
Men who have sex with men																	

■ Additional data


Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
Blood donors																	

Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
Tuberculosis patients	Major Urban Areas	N-sites		1	1		1		1					1			
		Minimum		33.3	40.1		56.5		59.7					74			
		Median		33.3	40.1		56.5		59.7					74			
	Maximum		33.3	40.1		56.5		59.7					74				
	Outside Major Urban Areas	N-sites			1			1				1					
		Minimum			41			64.8				84.3					
Median				41			64.8				84.3						
Maximum			41			64.8				84.3							



Maps of HIV sentinel sites

Mapping the geographical distribution of HIV sentinel sites for different population groups may assist in interpreting both the national coverage of the HIV surveillance system as well in explaining differences in levels and trends of prevalence. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, in collaboration with the WHO Public Health Mapping Team, Communicable Diseases, is producing maps showing the location and HIV prevalence of HIV sentinel sites in relation to population density, major urban areas and communication routes.

Trends in antenatal sentinel surveillance for higher prevalence countries, or in prevalence among selected populations for countries with concentrated epidemics, are a new addition. These will be presented for those countries where sufficient data exist.



Trends in HIV prevalence among antenatal clinic attendees



Median prevalence and ranges are shown in areas with more than one sentinel site.

The boundaries and names shown and the designations used on the map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.
 WHO 2002, all rights reserved.

6 - Zimbabwe

Reported AIDS cases

AIDS cases by year of reporting

1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
0	0	0	0	0	0	0	0	119	202	1311	4362	4557	8180	9174	10647	13356	12029	6732	4113		

2001 Total Unk

	74782	
--	-------	--

Date of last report: 30-Nov-1998

Following WHO and UNAIDS recommendations, AIDS case reporting is carried out in most countries. Data from individual AIDS cases are aggregated at the national level and sent to WHO. However, case reports come from surveillance systems of varying quality. Reporting rates vary substantially from country to country and low reporting rates are common in developing countries due to weaknesses in the health care and epidemiological systems. In addition, countries use different AIDS case definitions. A main disadvantage of AIDS case reporting is that it only provides information on transmission patterns and levels of infection approximately 5-10 years in the past, limiting its usefulness for monitoring recent HIV infections.

Despite these caveats, AIDS case reporting remains an important advocacy tool and is useful in estimating the burden of HIV-related morbidity as well as for short-term planning of health care services. AIDS case reports also provide information on the demographic and geographic characteristics of the affected population and on the relative importance of the various exposure risks. In some situations, AIDS reports can be used to estimate earlier HIV infection patterns using back-calculation. AIDS case reports and AIDS deaths have been dramatically reduced in industrialized countries with the introduction of HAART (Highly Active Anti-Retroviral Therapy).

AIDS cases by mode of transmission

Hetero: Heterosexual contacts.
 Homo/Bi: Homosexual contacts between men.
 IDU: Injecting drug use. This transmission category also includes cases in which other high-risk behaviours were reported, in addition to injection of drugs.
 Blood: Blood and blood products.
 Perinatal: Vertical transmission during pregnancy, birth or breastfeeding.
 NS: Not specified/unknown.

Sex	Trans. Group	<97	1997	1998	1999	2000	2001	Unkn.	Total	%	
All	All										
	Hetero										
	Homo/Bi										
	IDU										
	Blood										
	Perinatal										
	Other knowr										
	Unknown										
	Male	All									
		Hetero									
Homo/Bi											
IDU											
Blood											
Perinatal											
Other knowr											
Unknown											
Female		All									
		Hetero									
	Homo/Bi										
	IDU										
	Blood										
	Perinatal										
	Other knowr										
	Unknown										
	NS	All									
		Hetero									
Homo/Bi											
IDU											
Blood											
Perinatal											
Other knowr											
Unknown											

AIDS cases by age and sex

Sex	Age	<97	1997	1998	1999	2000	2001	Unkn.	Total	%	
All	All	25385	6732	4113					36230	100.0	
	0-4	3177	725	506					4408	12.2	
	5-14	561	206	177					944	2.6	
	15-19	361	107	50					518	1.4	
	20-29	6671	1608	901					9180	25.3	
	30-39	7922	2146	1319					11387	31.4	
	40-49	3928	1107	689					5724	15.8	
	50-59	625	128	61					814	2.2	
	60+	569	149	95					813	2.2	
	NS	1571	556	315					2442	6.7	
	Male	All	3575	2168						5743	100.0
		0-4	370	290						660	11.5
		5-14	111	91						202	3.5
15-19		27	6						33	0.6	
20-29		602	323						925	16.1	
30-39		1206	719						1925	33.5	
40-49		694	431						1125	19.6	
50-59		79	37						116	2.0	
60+		117	66						183	3.2	
NS		369	205						574	10.0	
Female		All	2988	1929						4917	100.0
		0-4	339	215						554	11.3
		5-14	87	86						173	3.5
	15-19	79	44						123	2.5	
	20-29	971	576						1547	31.5	
	30-39	897	598						1495	30.4	
	40-49	379	256						635	12.9	
	50-59	48	23						71	1.4	
	60+	29	29						58	1.2	
	NS	159	102						261	5.3	
	NS	All		16						16	100.0
		0-4		1						1	6.3
		5-14		0						0	0.0
15-19			0						0	0.0	
20-29			2						2	12.5	
30-39			2						2	12.5	
40-49			2						2	12.5	
50-59			1						1	6.3	
60+			0						0	0.0	
NS			8						8	50.0	

Curable Sexually Transmitted Infections (STIs)

The predominant mode of transmission of both HIV and other STIs is sexual intercourse. Measures for preventing sexual transmission of HIV and STI are the same, as are the target audiences for interventions. In addition, strong evidence supports several biological mechanisms through which STI facilitate HIV transmission by increasing both HIV infectiousness and HIV susceptibility. Also significant is the observation of a sharp decline in the concentration of HIV in genital secretions when the infection is treated. Monitoring trends in STI can provide valuable information on the sexual transmission of HIV as well as the impact of behavioural interventions, such as promotion of condom use.

Clinical services offering STI care are an important access point for people at high risk for both AIDS and STIs, not only for diagnosis and treatment but also for information and education. Therefore, control and prevention of STIs have been recognized as a major strategy in the prevention of HIV infection and ultimately AIDS. One of the cornerstones of STI control is adequate management of patients with symptomatic STIs. This includes diagnosis, treatment and individual health education and counselling on disease prevention and partner notification. Consequently, monitoring different components of STI control can also provide information on HIV prevention within a country.

■ Reported STI syndromes

Syndrome	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	Total	Unk
Urethral discharge														
Genital Ulcer														
Vaginal discharge														
Lower Abdominal Pain														
Neonatal conjunctivitis														

Date of last report:

■ Incidence of urethral discharge, men

Year	Area	Age Group	Rate	N=

Comments:

Sources:

■ Syphilis prevalence, women

Percent of blood samples taken from women aged 15-24 that test positive for syphilis during routine screening at selected antenatal clinics.

Year	Area	Age Group	Rate	N=

Comments:

Sources:

8 - Zimbabwe

Estimated size of populations at increased risk of HIV infection

	Year	Area	High estimate	Low estimate
Number of female sex workers				
Number of injecting drug users				
Number of men who have sex with men				

Comments:

Sources:

Health service and care indicators

HIV prevention strategies depend on the twin efforts of care and support for those living with HIV or AIDS, and targeted prevention for all people at risk or vulnerable to the infection. It is difficult to capture such a large range of activities with one or just a few indicators. However, a set of well-established health care indicators may help to identify general strengths and weaknesses of health systems. Specific indicators, such as access to testing and blood screening for HIV, help to measure the capacity of health services to respond to HIV/AIDS - related issues.

■ Access to health care

Indicators	Year	Estimate	Source
% of population with access to health services - total:			
% of population with access to health services - urban:			
% of population with access to health services - rural:			
Contraceptive prevalence rate (%):	1999	53.5	UNICEF/UNPOP
Percentage of contraceptive users using condoms:			
% of births attended by skilled health personnel:	1999	72.5	WHO
% of 1-yr-old children fully immunized - DPT:	2000	77	WHO/UNICEF
% of 1-yr-old children fully immunized - Measles:	2000	70	WHO/UNICEF
% of ANC clinics where HIV testing is available:			
% of PLWHA who have access to ARV:			

■ Number of people living with HIV/AIDS (PLWHA) receiving highly active antiretroviral therapy (HAART)

	1995	1996	1997	1998	1999	2000	2001	Total	Unk
People initiating HAART therapy									

■ Coverage of HIV Voluntary Counselling and Testing (VCT)

Number of functioning VCT sites per 100,000 population aged 15-49.

Year	Area	N=	Rate

Comments:

Sources:

Knowledge and behaviour

In most countries the HIV epidemic is driven by behaviours (e.g.: multiple sexual partners, injecting drug use) that expose individuals to the risk of infection. Information on knowledge and on the level and intensity of risk behaviour related to HIV/AIDS is essential in identifying populations most at risk for HIV infection and in better understanding the dynamics of the epidemic. It is also critical information in assessing changes over time as a result of prevention efforts. One of the main goals of the 2nd generation HIV surveillance systems is the promotion of a standard set of indicators defined in the National Guide (Source: National AIDS Programmes, A Guide to Monitoring and Evaluation, UNAIDS/00.17) and regular behavioural surveys in order to monitor trends in behaviours and to target interventions.

The indicators on knowledge and misconceptions are an important prerequisite for prevention programmes to focus on increasing people's knowledge about sexual transmission, and, to overcome the misconceptions that act as a disincentive to behaviour change. Indicators on sexual behaviour and the promotion of safer sexual behaviour are at the core of AIDS programmes, particularly with young people who are not yet sexually active or are embarking on their sexual lives, and who are more amenable to behavioural change than adults. Finally, higher risk male-male sex reports on unprotected anal intercourse, the highest risk behaviour for HIV among men who have sex with men.

■ **Knowledge of HIV prevention methods**

Proportion of people citing correctly at least two acceptable ways of protection from HIV infection.

Year	Area	Age Group	Male	Female	All
------	------	-----------	------	--------	-----

Comments:
Sources:

■ **Misconception about AIDS (no incorrect beliefs)**

Proportion of people who correctly reject the two most common local misconceptions about AIDS transmission or prevention, and who know that a healthy looking person can transmit AIDS

Year	Area	Age Group	Male	Female	All
------	------	-----------	------	--------	-----

Comments:
Sources:

■ **Median age at first sexual experience**

The age by which one half of young men or young women aged 15-24 have had penetrative sex (median age) of all young people surveyed.

Year	Area	Age Group	Male	Female	All
------	------	-----------	------	--------	-----

Comments:
Sources:

■ **Higher risk sex in the last year (adults)**

Proportion of adult respondents who have had sex with a non-regular (non-marital, non-cohabiting) partner in the last 12 months, of all adult respondents reporting sexual activity in the last 12 months.

Year	Area	Age Group	Male	Female	All
1999	All	15-49		16.0	
		15-59	42.5		

Comments: w = 15-59 age cohort
Sources: DHS

■ **Young people having multiple partners in last year (youth)**

Proportion of respondents who have had sex with more than one partner in the last 12 months.

Year	Area	Age Group	Male	Female	All
1999	All	15-24	10.8	2.0	

Comments:
Sources: DHS

Knowledge and behaviour

■ Condom use in last higher risk sex (adults)

The percentage of adult respondents who say they used a condom the last time they had sex with a non-regular (non-marital, non-cohabiting) partner, of those who have had sex with such a partner in the last 12 months.

Year	Area	Age Group	Male	Female	All
------	------	-----------	------	--------	-----

Comments:

Sources:

■ Young people using a condom during premarital sex (youth)

Proportion of young single people who used a condom at last sex.

Year	Area	Age Group	Male	Female	All
------	------	-----------	------	--------	-----

Comments:

Sources:

■ Commercial sex in the last year

Proportion of men reporting sex with a sex worker in the last 12 months.

Year	Area	Age Group	Rate	All
------	------	-----------	------	-----

Comments:

Sources:

■ Reported condom use in commercial sex

Proportion of men reporting condom use the last time they had sex with a sex worker, of those who report having had sex with a sex worker in the last 12 months.

Year	Area	Age Group	Rate	All
------	------	-----------	------	-----

Comments:

Sources:

■ Higher risk male-male sex in the last year

The percentage of men who have had anal sex with more than one male partner in the last 6 months, of all men surveyed who have had sex with a male partner.

Year	Area	Age Group	Rate	All
------	------	-----------	------	-----

Comments:

Sources:

■ Injecting drug users sharing equipment at last injection nationwide

Percentage of injecting drug users active in the last month who report sharing injecting equipment the last time they injected drugs.

Year	Area	Age Group	Rate	All
------	------	-----------	------	-----

Comments:

Sources:

Prevention Indicators

Male and female condoms are the only technology available that can prevent sexual transmission of HIV and other STIs. Persons exposing themselves to the risk of sexual transmission of HIV should have consistent access to high quality condoms. AIDS Programs implement activities to increase both availability of and access to condoms. These activities should be monitored and have resources directed to problem areas. The indicator below highlights the availability of condoms. However, even if condoms are widely available, this does not mean that individuals can or do access them.

■ **Condom availability nationwide**

Total number of condoms available for distribution nationwide during the preceding 12 months, divided by the total population aged 15-49.

Year	N	Rate
------	---	------

Comments:

Sources:

■ **Prevention of mother-to-child transmission (MTCT) nationwide**

Percentage of women who were counselled during antenatal care for their most recent pregnancy, accepted an offer of testing and received their test results, of all women who were pregnant at any time in the preceding two years.

Year	N	Rate
------	---	------

Comments:

Sources:

Blood safety programs aim to ensure that the majority of blood units are screened for HIV and other infectious agents. This indicator gives an idea of the overall percentage of blood units that have been screened to high enough standards that they can confidently be declared free of HIV.

■ **Screening of blood transfusions nationwide**

Percentage of blood units transfused in the last 12 months that have been adequately screened for HIV according to national or WHO guidelines.

Year	N	Rate
------	---	------

Comments:

Sources:

12 - Zimbabwe

Sources

Data presented in this Epidemiological Fact Sheet come from several different sources, including global, regional and country reports, published documents and articles, posters and presentations at international conferences, and estimates produced by UNAIDS, WHO and other United Nations agencies. This section contains a list of the more relevant sources used for the preparation of the Fact Sheet. Where available, it also lists selected national Web sites where additional information on HIV/AIDS and STI are presented and regularly updated. However, UNAIDS and WHO do not warrant that the information in these sites is complete and correct and shall not be liable whatsoever for any damages incurred as a result of their use.

Asamoah-Odei, E. 1999 Surveillance of HIV/AIDS and Related Diseases WHO Travel Report Summary, Zimbabwe, 12 July, unpublished trip report.

Armstrong, K., G. Woelk, S. Gloyd 2000 Socio-Economic Factors Related to HIV Prevalence among Antenatal Clinic Attendees at Sentinel Surveillance Sites Throughout ... XIII International AIDS Conference, Durban, South Africa, 7/9-14, Poster TuPeC3434.

Eriki, P. 1992 HIV Seroprevalence among Patients with Tuberculosis in Selected African Countries VII International Conference on AIDS in Africa, Yaounde, Cameroon, 12/8-11, Poster W.PS.Congo.

Easterbrook, P., D. Lamprecht, N. Ives, et al. 2000 Transmission Patterns of Tuberculosis in Harare, Zimbabwe: A Molecular Epidemiological Analysis XIII International AIDS Conference, Durban, South Africa, 7/9-14, Abstract WePeC4436.

Gwanzura, L., P. R. Mason, A. S. Latif, et al. 1996 Vaginal Lactobacilli and HIV Transmission in African Women XI International Conference on AIDS, Vancouver, 7/7-14, Abstract Th.C.4509.

Gregson, S., T. Zhuwau, R. M. Anderson, et al. 1995 Age and Religion Selection Biases in HIV-1 Prevalence Data from Antenatal Clinics in Manicaland, Zimbabwe Central African Journal of Medicine, vol. 41, no. 11, pp. 339-346.

Gregson, S., C. Nyamukapa, P. Mason, et al. 2000 Population-Based Survey of HIV Infection and Its Socio-Demographic and Behavioural Determinants in Rural Manicaland, ... XIII International AIDS Conference, Durban, South Africa, 7/9-14, Poster TuPeC3447.

Guevara, H., E. Johnston, L. Zijenah, et al. 2000 Prenatal Transmission of Subtype C HIV-1 in Zimbabwe: HIV-1 RNA and DNA in Maternal and Cord Blood Journal of Acquired Immune Deficiency Syndromes, vol. 25, no. 5, pp. 390-397.

Jackson, H. 1993 AIDS Update in Zimbabwe: AIDS Cases ZAINET AIDS News, vol. 1, no. 3, pp. 10-13.

Legg, W., M. Mahari, S. Houston, et al. 1989 Association of Tuberculosis and HIV Infection in Zimbabwe V International Conference on AIDS, Montreal, 6/4-9, Abstract Th.G.O. 6.

Le Bacq, F., P.R. Mason, L. Gwanzura, et al. 1993 HIV and Other Sexually Transmitted Diseases at a Rural Hospital in Zimbabwe Genitourinary Medicine, vol. 69, pp. 352-256.

Latif, A. S. 1995 A Report on a Study to Determine the Aetiology and Pattern of STD amongst Men and Women Presenting to Health Centres in ... University of Zimbabwe Medical School, Dept. of Medicine, September STD report, unpublished.

Mertens, T., G. Tondorf, M. Siebolds, et al. 1989 Epidemiology of HIV and Hepatitis B Virus (HBV) in Selected African and Asian Populations Infection, vol. 17, no. 1, pp. 4-7.

Mahomed, K., J. Kasule, D. Makuyana, et al. 1991 Seroprevalence of HIV Infection amongst Antenatal Women in Greater Harare, Zimbabwe Central African Journal of Medicine, vol. 37, no. 10, pp. 322-325.

Mahari, M., W. Legg, S. Houston, et al. 1990 Association of Tuberculosis and HIV Infection in Zimbabwe VI International Conference on AIDS, San Francisco, 6/20-24, Abstract Th.B.494.

Mbizvo, M. T., A. Mashu, T. Chipato, et al. 1996 Trends in HIV-1 and HIV-2 Prevalence and Risk Factors in Pregnant Women in Harare, Zimbabwe Central African Journal of Medicine, vol. 42, no. 1, pp. 14-21.

Mason, P., S. Ray, C. Mapushere, et al. 1995 Use of Female and Male Condoms by Commercial Sex Workers: Impact on STD Transmission IX International Conference on AIDS and STD in Africa, Kampala, Uganda, 12/10-14, Poster TuC614.

Olayinka, B. A., C. L. OBI 1999 Symptomatic HIV-Infection in Infants According to Serostatus of Mothers during Pregnancy East African Medical Journal, vol. 76, no. 10, pp. 566-570.

Schoch, O. D., H. L. Rieder 1996 Characteristics of Sputum Smear-Positive Tuberculosis Patients with and without HIV Infection in a Hospital in Zimbabwe European Respiratory Journal, vol. 9, pp. 284-287.

Tswana, S. A., L. Nystrom, S. R. Moyo, et al. 1995 Hospital-Based Study of Sexually Transmitted Diseases at Murewa Rural District Hospital, Zimbabwe 1991-1992 Sexually Transmitted Diseases, vol. 22, no. 1, pp. 1-6.

Whiteside, A. 1991 HIV Infection and AIDS in Zimbabwe: An Assessment Southern Africa Foundation for Economic Research, Economic Research Unit, University of Natal, pp. 1-50.

Zimbabwe Ministry of Health 1990 HIV Sentinel Surveillance 1990 Report AIDS Control Programme and Health

Information Unit, Ministry of Health, report.

Zimbabwe Ministry of Health 1991 HIV and AIDS Surveillance Quarterly Report : January, February, March 1991
AIDS Control Programme and Health Information Unit, Ministry of Health, report.

Zhuwau, T. 1997 HIV Trends and Behaviour Changes Presented at the UNAIDS Regional Workshop on "Evidence of Behavioural Change in the Context of HIV Decline in Uganda." 10-13 February, Nairobi, Kenya.

Zimbabwe Ministry of Health and Child Welfare 1994 HIV, STD, and AIDS Surveillance Workshop: Monomotapa Hotel
Health Information Unit, Department of Epidemiology and Disease Control, Ministry of Health and Child Welfare, 7-8
July, report.

Zimbabwe Ministry of Health and Child Welfare 1997 Summary Results of HIV Sentinel Surveillance AIDS
Coordination Programme, fax from Sunday Manyenya, 21 April.

Zimbabwe Ministry of Health and Child Welfare 1998 HIV, AIDS, STD and TB Fact Sheet Monitoring and Evaluation
Unit, National AIDS Coordination Programme, November, report, pp. 1-18.

Zimbabwe Ministry of Health 1998 HIV Sentinel Surveillance Data Ministry of Health & CW EDC-NHIS Unit,
documents.

Zimbabwe Ministry of Health and Child Welfare 2000 National Survey of HIV and Syphilis Prevalence among Women
at Antenatal Clinics in Zimbabwe, 2000 Republic of Zimbabwe, Ministry of Health and Child Welfare, CDC AIDS
Project, unpublished report.

Websites: www.aids.africa.com

13 - Zimbabwe

Annex: HIV Surveillance by site

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001			
Pregnant women	Major Urban Areas	Bulawayo			10.00		17.10			25.80		30.00		24.00			31.10		
		Chitungwiza								41.70									
		Chitungwiza (Seke North Clinic)												32.00					
		Chitungwiza (St. Mary Clinic)												34.00				33.50	
		Harare				23.80									28.00			30.00	
		Harare (6 peri-urban clinics)									32.00								
		Harare (Budiro)											30.40						
		Harare (Edith Opperman Clinic)				17.90													
		Harare (Glenview Clinic)				19.40													
		Harare (Greater Harare)									30.40				22.70				
		Harare (Harare Hospital)				16.00									33.33				
		Outside Major Urban Areas	Banket						15.70										27.80
			Beitbridge								23.00	44.00	59.00	46.00	37.00				41.40
			Bindura								40.00	27.00			29.30				32.10
			Binga (Binga Hosp.)												9.20				
	Binga (Binga Kariyangwe)																	13.00	
	Binga district (Binga Hosp.)								17.50		14.40								
	Binga district (Kariyangwe Hosp.)								6.60										
	Birchenough							15.50	24.00										
	Buhera													50.80					
	Chipinge									13.70									
	Chiredzi							39.70				70.20	46.70	25.00				70.70	
	Gokwe district (Growth Point)								22.00	27.00	34.00								
	Gokwe district (Midlands Pr.)							22.40											
	Guto										20.50	39.50		25.00				27.40	
	Gwanda						16.00	21.20	25.00	25.00	33.00	34.00	48.00	31.00				30.40	
	Gweru						28.10	24.00			34.50								
	Gweru (Mkoba 1)														19.20				
	Gweru (Mkoba Polyclinic)														30.70			36.40	
	Gweru (private clinic)														24.00				
	Honde Valley										24.30								
	Hwange (Wankie Colliery)										22.90								
	Hwange (Hwange Hosp.)					12.30	24.50	19.90						18.80					
Hwange (St. Patricks)																	32.60		
Hwange (Wankie Colliery Comp. Hosp.)							11.10												
Karanda									20.00				23.90				18.30		
Kariyangwe													7.00						
Kezi										16.00									
Kwekwe									25.00										
Kwekwe district (Mbizo 11 Clinic)													7.00						
Makoni														29.30					
Mandava						16.00													
Manicaland Pr.(Eastern Highlands)									15.80										
Maphisa										15.00			27.60	29.00					
Mary Mount													34.90						
Mashoko									19.10	18.10	19.00		30.00						
Masvingo						30.90	42.10			35.20		36.50							
Masvingo Province											41.70								
Matabeleland South Pr.(Antelope Hosp.)										15.00									
Matabeleland South Pr.(Antelope Hosp.)												28.00							
Mberengwa district (Mneme & Musume)					7.60														
Mberengwa district (Mneme & Musume)						7.70	7.70		25.50			30.00							
Mberengwa district (Musume Mission Hosp.)																54.50			
Midlands Pr. (commercial farms)								22.50	36.20										
Midlands Prov. (Mines)									24.50										
Murambinda				19.70	13.90					41.40						25.70			
Mutare								25.20		33.60		37.70							
Mutasa & rural areas															20.80				
Mutoko						25.90			34.40			19.10				33.70			
Mutorashanga						15.00													
Mvurwi												27.00							
Nkulumane																31.10			
Plumtree					9.90	11.10													
Rusape				31.60	33.80					67.00		53.30							
Rusitu Valley									14.00										
Sadza												31.40				42.70			
Sakubva																33.90			
Sanyati						20.00	18.50	18.50								36.90			

						17.00														
						23.70	20.00	17.00												
										23.00		23.00								
											37.30									
											42.60				55.70					
											24.30									
											44.00									
Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001			
Sex workers	Major Urban Areas	Harare									86.00									
Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001			
Injecting drug users																				
Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001			
STI patients, Males/both & females	Major Urban Areas	B/ Bulawayo					39.00		60.30		71.00									
		B/ Chitungwiza (St. Mary's Clinic)						70.00												
		B/ Chitungwiza (Zengeza Clinic)						56.00												
		B/ Harare				52.00	51.00													
		Harare								50.70	53.00									
		M/ Harare								71.20										
	Outside Major Urban Areas	B/ Banket/rural						42.40												
		B/ Beitbridge/rural								73.00	65.00									
		B/ Bindura								56.00	43.00									
		B/ Binga/rural						44.60												
		B/ Gokwe dist.				48.80	49.40	32.20												
		B/ Gutu/rural							71.80											
		B/ Gwanda/rural					33.10	40.00		66.00	66.00									
		B/ Gwenu					48.20	48.00	52.00	52.00										
		B/ Hwange (Hwange Hospital)				28.70	42.00	42.30												
		B/ Hwange (Wankie Colliery Hosp.)					32.60													
		B/ Kwekwe dist./rural							55.70											
		B/ Kwekwe/rural					45.60	48.80												
		B/ Mashoko/rural								51.90	58.40									
		B/ Mashonaland West Province							52.10											
		B/ Masvingo					58.50	59.50		75.40		71.80								
		B/ Matebeleland North Pro.				32.60														
		B/ Mberengwa dist./rural				24.30	24.49		46.80											
		B/ Midlands Prov. (Commercial Farms)								51.00										
		B/ Midlands Prov. (Mines)								48.00										
		B/ Murambinda/rural									69.30									
		B/ Mutare							25.60		56.60									
		B/ Mutorashanga/rural						39.00												
		B/ Plumtree/rural					33.30	43.10												
		B/ Rusape				59.70	56.40				87.70									
		B/ Sanyati					45.60		62.20											
		B/ Shabanie					46.70													
		B/ Shurugwi					39.60	43.00	41.00											
		F/ Karoi dist./rural					65.30													
		F/ Murewa dist.						46.00												
		F/ Mutoko/rural						43.60												
		M/ Karoi dist./rural					60.60													
		M/ Murewa dist.						46.00												
		M/ Mutoko/rural						59.60												
		M/ Rural area					6.60													
		Rural area					5.20													
Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001			
Men having sex with men																				

Additional data

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
Blood donors																
Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
Tuberculosis patients	Major Urban Areas					56.50		59.70								
			33.30	40.10								74.00				
	Outside Major Urban Areas			41.00		64.80										
									84.30							